






Copyright Pearson Prentice Hall


20-2 Animal-like Protists: Protozoans





Structures found in sarcodines that are used for feeding and movement are known as

- a. pseudopods.
- b. flagella.
- c. cilia.
- d. food vacuoles.





Structures found in sarcodines that are used for feeding and movement are known as

a. pseudopods.

- b. flagella.
- c. cilia.
- d. food vacuoles.





The structure found in most ciliates that contains a "reserve copy" of all the cell's genes is the

- a. macronucleus.
- b. micronucleus.
- c. trichocysts.
- d. contractile vacuole.





The structure found in most ciliates that contains a "reserve copy" of all the cell's genes is the

a. macronucleus.

- b. micronucleus.
- c. trichocysts.
- d. contractile vacuole.





- One way to classify the various groups of animal-like protists is by
 - a. the presence of a nuclear membrane.
 - b. the presence of mitochondria.
 - c. their means of movement.
 - d. the number of contractile vacuoles.



Slide 7 of 50

End Show

- One way to classify the various groups of animal-like protists is by
 - a. the presence of a nuclear membrane.
 - b. the presence of mitochondria.

c. their means of movement.

d. the number of contractile vacuoles.





4 Malaria is caused by the sporozoan

- a. Plasmodium.
- b. Anopheles.
- c. Amoeba.
- d. Paramecium.







Malaria is caused by the sporozoan

a. Plasmodium.

- b. Anopheles.
- c. Amoeba.
- d. Paramecium.



